International Journal of Psycho-Oncology and Positive Psychology

Vol (1) Issue (1) December 2023 ISSN− 2584-2439

Article

Antenatal and Postnatal Mental Health Care: Need of the Hour

Nandika Agarwal[1], Sneha Rustagi[1], Khushi Gupta[1], and Dr Smriti Sethi[2]

[1]Student, Amity Institute of Psychology and Allied Sciences, Amity University Uttar Pradesh, Noida, Uttar Pradesh 201301, India

[2] Assistant Professor, Amity Institute of Psychology and Allied Sciences, Amity University Uttar Pradesh, Noida, Uttar Pradesh 201301, India

Keywords:

Antenatal, Interventions, Maternal Well-Being, Mental Health, Postnatal, and Risk Factors.

Abstract

The transition from pregnancy to early motherhood is a critical stage that is characterized by significant physiological and psychological changes. The research aims to study the effect of antenatal and postnatal period on mental health of women. The purpose is to shed light on the prevalence of mental health issues, identify risk factors, and suggest appropriate interventions to lessen the associated burden through a thorough investigation. For achieving the objectives, an extensive review of literature was done to look at the body of knowledge on antenatal and postnatal mental health. To fully grasp the insights of the topic, 30 reviews of the literature were studied, covering a range of demographics, cultural situations, and mental health outcomes were examined. The analysis of the review papers made it possible to pinpoint common risk factors for mothers' mental discomfort. The discussion section emphasized on how complex pregnancy and postnatal mental health issues can be, including changes in hormone levels, social pressures, can and a lack of suitable support networks. The research also explores on how untreated maternal mental anguish has negative effects on not just the mother's well-being but also the growth of their children and the dynamics of the entire family. The essential need for a comprehensive strategy to address antenatal and postnatal mental health concerns is offered with the several interventions, from easily available mental health treatments to community support programs. A holistic support framework is promoted because of the significant impact on maternal well-being, child development, and societal fabric. Society can lessen the negative consequences and create a foundation for future generations of healthy individuals by including mental health awareness in prenatal and postpartum care.

Introduction

Antenatal and postnatal mental health refers to the psychological well-being of expecting mothers and new mothers throughout the first year following childbirth. The complex physiological, psychological, and emotional changes that take place throughout these critical phases have an impact not just on the mothers' mental health but also on the general well-being of their children and families. For fostering the healthy development of the mother and the child, it is crucial to comprehend the importance of mental health during these times.

Antenatal Period

Medical personnel actively monitor the pregnant woman's health during the antenatal phase; provide important knowledge and instruction about pregnancy, delivery, including postnatal care, and give necessary medical interventions to address any difficulties. This encompassing strategy strives to promote the normal growth of the fetus and improve mother health, eventually assuring a safe and healthy delivery. The antenatal period refers to the time during pregnancy before childbirth. According to World Health Organisation (WHO) (2020), The term "antenatal period" refers to the time during pregnancy, from conception to the start of labour or childbirth.

"The preconception and intrapartum periods are included in the definition of the antenatal period, which spans the period from conception until the end of pregnancy" as mentioned by F. Gary Cunningham in his book Williams Obstetrics (2018). Terri Kyle and Susan Scott Ricci's in 2019 expressed in their book titled 'Maternity and Paediatric Nursing' that the antenatal period, or prenatal period, encompasses the period from conception through the end of pregnancy, which is typically defined as childbirth. Also according to Murray Enkin and colleagues' in their book titled 'A Guide to Effective Care in Pregnancy and Childbirth', published in 1989 "The time from conception until the start of labour, which ends in childbirth, is referred to as the "antenatal period," also known as the "prenatal period."

Stages of Antenatal Period

First Trimester (Weeks 1-12). The egg that is fertilized implants in the uterus during conception. The embryo develops as a result of rapid cell division. Major organs and systems of the body start to form during a process called organogenesis. The placenta develops and begins supplying the embryo with nourishment and oxygen. Morning sickness, exhaustion, breast pain, and frequent urination are typical symptoms.

Second Trimester (Weeks 13-27). The fetus can more easily be identified as a human by the completion of the second trimester. The pregnant woman may feel the fetal movement as it intensifies. Organs keep growing and maturing. Ultrasound is frequently used to identify the gender of the fetus. The pregnant woman may feel less sick and have more energy. A developing belly and weight increase are more apparent

Third Trimester (Weeks 28-40+). The fetus continues to grow and weighs a substantial amount.Systems and organs are improved and made ready for life outside the womb.In order to give birth, the fetus settles in a head-down posture.Due to the uterusexpanding size, the pregnant woman may have increasing pain, backaches, frequent urination, and shortness of breath. "Practice contractions," also known as Braxton Hicks contractions, might become more obvious. As the due date draws near, the expecting mother may feel a variety of emotions and eagerness for labor and delivery.

Although these trimesters serve as a broad reference, it's crucial to remember that every woman's pregnancy experience is unique. The antenatal period is a special and transformational time for individuals who are expecting since the phases of pregnancy are marked by both physical modifications in the human body and emotional changes in the expectant person. A healthy antenatal time includes regular prenatal care, good diet, exercise, and emotional well-being.

Theoretical Basis to Understand the Antenatal Period

The theory underlying how antenatal period affects mental health of women is based on several psychological models and hypotheses. Here are a few significant theoretical stances that advance our comprehension of this phenomenon:

Antenatal Attachment Theory (John Bowlby and Mary Ainsworth, 1991). This theory focuses on the emotional connection that develops between the expectant mother and the growing fetus. It implies that a mother's emotions and attachment during pregnancy might affect how her unborn child develops emotionally and socially.

Maternal-Fetal Conflict Theory (Karl max). In this theory, possible conflicts of interest between a pregnant woman and her developing fetus are taken into account. It examines how a misalignment between resources and physiological requirements could occasionally lead to particular adaptations or responses.

Stress and Antenatal Development (Janet Di Pietro and Curt Sandman). Researchers investigate how mother stress during pregnancy affects fetal growth and long-term results for the kid. This involves being aware of how stress hormones may impact the developing fetus and may cause behavioral and developmental problems.

Nutritional Impact on Fetal Development (Barker). This idea emphasizes the crucial contribution of maternal nutrition to the appropriate growth, organ development, and general health of the fetus during pregnancy.

Maternal Bonding and Mental Health (Bowlby). Both maternal and fetal outcomes can be significantly influenced by the emotional and mental health of the pregnant person. Systems of support and constructive mental health behaviors are essential.

Causal Factors Affecting Women's Mental Health During Antenatal Period

Social Support. A female's mental health during pregnancy can be significantly improved by having a solid support network of family, friends, and a partner. Stress and anxiety can be exacerbated by a lack of support or a sense of isolation.

Financial Stress. Anxiety and stress during pregnancy can be exacerbated by financial uncertainty or concerns about meeting the requirements of the unborn child.

Cultural Expectations. A woman's mental health may be impacted by social norms and standards around pregnancy and parenthood. She can have low self-esteem and inadequacy as a result of these expectations.

Job-Life Balance. Juggling a pregnancy and job obligations can be difficult. Negative mental health impacts may occur in women who endure workplace discrimination, a lack of paid maternity leave, or difficulties connected to their jobs.

Previous Mental Health History. Pregnant women with a history of mental health problems are more likely to encounter difficulties. Existing problems may become worse as a result of hormonal changes and lifestyle changes.

Healthcare Access. A lack of access to high-quality healthcare and prenatal services might result in doubt about the health of the pregnancy, which may cause stress and worry.

Traumatic Experiences. Past traumatic events, such as abuse or bereavement, might return during pregnancy and have an impact on a woman's mental health. Promoting the good mental health of pregnant women can be significantly aided by addressing these issues and offering the right assistance.

Challenges Faced by Women During Antenatal Period

Emotional Changes and Mood Swings. Hormonal changes that frequently accompany pregnancy can cause mood swings, impatience, and emotional sensitivity. Neurotransmitter level changes can affect how well mood management works.

Anxiety and Stress. Stress levels can be increased by anxiety related to a baby's health, childbirth, and parenthood. Stress can be exacerbated by alterations in lifestyle and financial problems.

Depression and Perinatal Mood Disorders. Antenatal depression, or depression throughout pregnancy, is a possibility. Depression and anxiety are considered perinatal mood disorders throughout pregnancy and after delivery. Also, Changes in body composition and weight increase can have an impact on one's self-esteem, and societal standards and conceptions of pregnancy may also play a role.

Maternal-Fetal Attachment. While some women have a deep emotional bond with their unborn child, others may find it difficult. Guilt or anxiety from not feeling attached might have an impact on one's mental health.

Support Networks and Social Isolation. Having a strong network of friends, family, and partners can help your mental health. Stress and feelings of separation from others can be caused by social isolation or a lack of support.

Postnatal Period

The term "postnatal" describes the period of time right after delivery. It is the stage that follows childbirth and lasts during the first few weeks or months of a child's existence. Given that it entails major mental, physical, and psychological changes and adaptations for both the mother and the infant, this time period is crucial for both.

The World Health Organization (WHO) defined "postnatal" in 2004 as the time between delivery and the first six weeks of life when the infant experiences a variety of physiological and behavioral adjustments.

The term "postnatal" refers to the period of time immediately following childbirth, which includes the mother's physical and mental rehabilitation as well as the care of the infant, according to the American College of Obstetricians and Gynaecologists.

According to Adele Pillitteri's textbook "Maternal and Child Health Nursing: Care of the Childbearing and Childrearing Family," "postnatal" is the time following birth when the

mother's body starts to recover to its pre-pregnancy form and the infant adjusts to life outside the womb.

"Postnatal" is defined by the Merriam-Webster Medical Dictionary as the time following childbirth, which includes both the mother's recuperation and the development of the infant.

Stages of Postnatal Period

Immediate Postpartum Stage (First 24 Hours). The immediacy of postpartum is shown in this stage. Medical experts constantly watch the mother and child throughout this period for any serious health issues. The primary goals are to manage any potential difficulties, stabilize the mother's condition, and see to the baby's fundamental necessities.

Early Postpartum period (First Week*).* Starting the second day following labor, this period lasts through the first week. During this phase, the mother's body experiences major physical and hormonal changes. Uterine contractions, vaginal discharge (lochia), breast engorgement when milk production starts, and emotional changes as the mother adjusts to her new role are typical occurrences. During this phase, medical examinations may take place to evaluate the mother's recovery and address any issues.

Late Postpartum Stage (Six Weeks*).* This phase begins at the conclusion of the first week following delivery and lasts for six weeks. The mother's body is still healing and returning to its pre-pregnancy form at this period. The woman may endure physical discomfort, exhaustion, and emotional fluctuations as her uterus progressively shrinks to its usual size. The mother should gradually resume her regular activities and seek medical help if any difficulties develop.

Remember that every woman's postnatal experience is unique, and the stages may overlap or differ in length. Additionally, seeking the advice of medical specialists for precise and individualized advice throughout the postnatal time is always advised.

Theoretical Basis to Understand the Postnatal Period

The theory underlying how postnatal period affects mental health of a women is based on several psychological models and hypotheses. Here are a few significant theoretical stances that advance our comprehension of this phenomenon:

The Idea of Parental Role Development (Baumrind, 1966). According to this idea, people go through many phases and changes when they become parents, including after giving birth. It looks at how parents adjust to their new roles, attitudes, and actions with regard to the newborn.

Social Support Theory (Francis Cullen, 1994). In times of change, such as the postnatal period, this theory highlights the value of social networks and support systems. It emphasizes how social support may reduce the strains and difficulties of being a new parent.

Transactional Model of Stress and Coping (Lazarus and Folkman, 1987). According to this model, both internal and environmental variables can have an impact on how much stress a person experiences after giving birth. It looks at how people experience and manage stress and how it affects their mental health.

Attachment Theory (Bowlby, 1958). This theory examines the emotional ties that develop between parents and their young children. It is pertinent to the postnatal period because it

looks at how the parent-child bond affects both the development of the infant and the well-being of both parents.

Transition to Parenthood Theory (Raphael-Leff, 2001 and Stern and Bruschweiler-Stern, 1998). In this theory, the psychological, emotional, and relationship changes that people and couples go through when they have children are highlighted. It explores the difficulties and adjustments during the postnatal period.

Causal Factors Affecting Women's Mental Health During Postnatal Period

Maternal Role Adjustment. Getting used to motherhood's obligations might be difficult. Having too many duties or feeling inadequate might affect a woman's mental health.

Sleep Deprivation. Disrupted sleep patterns brought on by caring for a baby can lead to exhaustion, irritability, and mood swings, which can have an impact on mental health.

Body Image Issues. Changes in a woman's body image after giving birth might lower her self-esteem and make her feel anxious and depressed.

Employment-Life Balance. For new mothers, juggling employment, obligations to one's family, and self-care may be difficult. Mental health may be impacted by workplace support, pregnancy leave rules, and flexible work schedules.

Previous Mental Health History. Women who have experienced mental health problems in the past may be more vulnerable to postpartum mental health concerns.

Challenges Faced by Women During Postnatal Period

Postnatal depression. Signs include persistent melancholy, loss of interest, changes in eating and sleep patterns, exhaustion, and feelings of worthlessness. American College of Obstetricians and Gynaecologists is a good source.

Postnatal anxiety. Signs and symptoms include excessive concern, restlessness, irritation, trouble focusing, and bodily manifestations like shaking or sweating.Postpartum Support International as a source

Postnatal PTSD. Signs and symptoms include nightmares, avoiding triggers, increased arousal, and emotional numbness. Postpartum Support International as a source

Postnatal Bipolar Disorder. Changes in sleep, activity levels, and behavior, as well as periods of significant mood swings, ranging from manic (elevated mood) to depressed episodes. National Institute of Mental Health as a source

Breastfeeding difficulties. Nipples that are painful or cracked, engorgement, trouble latching, a lack of milk production, or mastitis (breast infection) are all symptoms.

Body Image Issues. Feeling self-conscious about postpartum physical changes, gaining weight, getting stretch marks, or body shape changes.

It's crucial for anyone dealing with any of these problems to get help and assistance from medical specialists. Pregnancy-related mental health problems are equally as important as pregnancy-related physical health problems, and earlier intervention can benefit both the mom and the unborn child.

Literature Review

Deepthi et al. (2023) conducted a study on ‘The Acceptability of a mHealth App for monitoring perinatal and postpartum mental health: A qualitative study with women and professionals’. The study's conclusions demonstrate that pregnant and postpartum women would embrace the use of mental health to track their mood problems. This could help inform the creation of low-cost, therapeutically useful instruments that make it easier to continuously monitor, identify, and treat mood disorders in this vulnerable population.

Kodi et al. (2023) conducted a study on ‘The COVID-19 pandemic's social determinants of prenatal mental health’. According to the findings, social resources have the strongest correlation with mental health, making them noteworthy unique correlations. Although no moderating effects of composite resource measures were discovered, BIPOC child bearers showed a greater correlation between pandemic employment-related lowered resources and symptoms than White child bearers did. Thus, it was determined that prenatal mental health is influenced by both consistent social-structural variables and acute crisis-related alterations, with larger levels and/or impacts of resources helping to explain racial and ethnic inequalities.

Jennifer et al. (2022) conducted a study on a ‘New hazard to perinatal mental health climate change’. The conclusion points to a period that may be difficult for women for a number of reasons. A few of the elements that can affect perinatal mental health include necessary role adjustments (reprioritization), changes in one's capacity to access pre-birth levels (and types) of social support, fluctuating hormones, changes in body shape, and potential complications during pregnancy, childbirth, or postpartum. As a result, it is a crucial omission because the mother's mental health has an effect on the entire family.

Krystal (2022) conducted a study on ‘Employment and maternal (perinatal) mental health: A research and practical agenda’. The findings imply that scientists should take into account the interaction of four factors: the body, the sociocultural milieu, personal agency, and time. At various levels of a framework for bio-ecological systems, a number of concerns for research are suggested. These issues are based on current transdisciplinary literature and illustrative examples from the UK setting.

Roxanne et al. (2022) conducted a study on a ‘Qualitative interview study on the views of pregnant women and healthcare professionals on perinatal mental health and intimate partner abuse in rural Ethiopia’. According to the research, women and health professionals in rural Ethiopia believe that there are several interrelated paths linking IPV and prenatal mental problems. Contrary to assumptions of tact, women and healthcare professionals felt comfortable talking about the effect of IPV on perinatal mental health and endorsed the necessity for quick mental health interventions integrated into ANC.

Theano and Eleftheria (2022) conducted a study on ‘Restrictions connected to the COVID-19 pandemic: Elements that may have an impact on perinatal maternal mental health and possible repercussions for newborn development’. According to the research, the pandemic may have had a negative impact on pregnant women's mental health, which could have a negative impact on babies from the Covid-19 generation. In an effort to support maternal mental health and newborn development, they emphasize the need for evidence-based therapies to be integrated throughout the healthcare system for prenatal and postpartum care.

Vikki et al. (2022) conducted a study on ‘Why, how, and what's next: Perinatal mental health education program for maternity, neonatal, and other allied multidisciplinary teams in NHS lothian’. The findings indicated that the sessions were understandable to the participants and that the content was clinically helpful and pitched appropriately. Despite COVID-19's difficulties when the program first began, participants have praised how collaborative and supportive the sessions have been delivered. Understanding what participants are learning and how it is affecting their practice will be the next step in the development of the program.

Zahra et al. (2022) conducted a study on ‘Findings from the norwegian mother, father, and child cohort study showing a link between prenatal work stress and prenatal and postnatal depression and anxiety’. According to the findings, pregnant women who are under stress at work are more likely to later suffer from melancholy and anxiety. The findings can guide the creation of workplace policies to help new and expectant moms' mental health.

Billie et al. (2021) conducted a study on ‘A qualitative investigation of models of community mental health care for women with perinatal mental health problems’. The findings suggest that women who experience perinatal mental health difficulties value specialist perinatal expertise, but that general, non-perinatal teams may also have advantages for some. Further research into optimal care arrangements is merited.

Signe et al. (2021) conducted a study on ‘Parental mental health conditions and the use of healthcare services in children during the first year of life– a register-based, nationwide study’. The results suggest that both maternal and paternal mental health conditions, including minor mental health conditions, were associated with increased utilization of healthcare services. Focus on both parents’ mental health conditions (even if minor) may be warranted in service planning.

Methodology

Aim

To Study the effect of Antenatal and Postnatal Period on Women’s Mental Health

Objectives

- To study the underlying factors and effects of antenatal and postnatal period on mental health of women
- To review the researches on the effects of antenatal and postnatal period on mental health of women
- To recommend strategies for promotion of positive mental health of women during the antenatal and postnatal period.

Procedure

A secondary research review of 30 papers was done.

Discussion

The immediate need for antenatal and postnatal mental health care cannot be denied. These crucial times span the emotional health of pregnant mothers during pregnancy (antenatal) and the months immediately following childbirth (postnatal), which are

sometimes overlooked in terms of their importance. It is crucial to acknowledge the significance of comprehensively addressing these concerns for a number of reasons.

Maternal mental health has a significant impact on both the mother and child. The mother's mental state is extremely important during the vital prenatal time for fetal development. Preterm birth and low birth weight are only a couple of the negative effects that stress, anxiety, and depression during pregnancy have been related to. Additionally, postpartum mental health problems might make it more difficult for mothers to care for their children and build emotionally close relationships with them. For safe pregnancies and the best possible development of the child, it is crucial to give antenatal and postnatal mental health a top priority.

The effects of maternal mental health extend far beyond the immediate family. A mother's mental health, including her partner and other family members, greatly impacts the overall dynamics of a family. Untreated mental health problems can cause family conflict and interpersonal tension, which may have an adverse effect on the child's long-term emotional growth. By addressing antenatal and postnatal mental health, families can become healthier and children can have stronger foundations.

In order to lessen the ubiquitous stigma associated with mental health disorders, particularly in the context of motherhood, it is imperative to address antenatal and postnatal mental health. When mental health issues occur, there is a damaging cultural expectation that moms should always be happy and pleased. This expectation can cause mothers to feel ashamed and alone. Fighting this stigma, encouraging women to seek assistance when necessary, and developing a more compassionate and inclusive society may all be accomplished by openly discussing these difficulties and offering proper support.

Furthermore, it's critical to understand that being a mother is a huge life event that frequently results in considerable physical, emotional, and social changes. Because of how stressful this change can be, women are more likely to experience mental health problems. Therefore, providing comprehensive mental health assistance to expectant and new mothers can aid women in navigating this profound change in their lives more successfully while lowering the likelihood of mental health issues and their long-term effects.

It should be noted that antenatal and postnatal mental health support is an urgent necessity rather than merely a suggestion. Beyond the individual, it has an effect on the family and society as a whole. In order to encourage healthier pregnancies, stronger families, and a more compassionate society, we have to place a high priority on the emotional health of mothers both during and after pregnancy. A critical first step in guaranteeing mothers' well-being and the future of our society is realizing that mental health issues at these times are both frequent and treatable. It's time to accept that taking care of antenatal and postnatal mental health is essential for a community to be healthier, more compassionate, and more resilient.

Pouran et al. (2023) conducted a study on ‘Content analysis research on maternal mental health issues during the COVID-19 pandemic in Iran’. The findings imply that the primary worry and dread of pregnant and postpartum women during the COVID-19 epidemic was the risk of their own or their unborn child's mortality. Managers should plan to strengthen and encourage women's mental health, especially in stressful conditions, with the knowledge that was learned by understanding pregnant women and new moms about mental health concerns during the COVID-19 pandemic.

Amelia and Carol (2022) conducted a study on ‘The needs and experiences of New Zealand mothers and healthcare professionals with regard to postpartum depression and anxiety during the COVID-19 pandemic’. The findings of this study point to the necessity of giving postnatal women safe, on-site access to healthcare practitioners and sources of social support during pandemic lockdowns in order to lessen isolation at this delicate time. Expanding the range of support options available to mothers, both inside and beyond the context of the epidemic, may be facilitated by establishing a dedicated postnatal mental health support line. It is also advised that midwives and other postnatal healthcare professionals, such as well child nurses, focus more on mental health training to improve their capacity to assist women who are experiencing postnatal mental illness.

Céline and Jaqueline (2022) conducted a study on ‘The Impact on mothers' postnatal sense of security and mother-to-infant attachment during the COVID-19 epidemic’. The results emphasize the significance of taking social and environmental requirements and factors into account when assessing postnatal mental health and offering postnatal care to new moms during a medical emergency. In order to prevent parental isolation during the delicate postpartum period, health services and experts should pay particular attention to mothers' mental health and well-being and ensure continuity of treatment.

Katie et al. (2021) conducted a study on’ How music may support perinatal mental health' an overview’. The result includes single studies and reviews that use both qualitative and quantitative methods. Drawing on animal and human models, the effect of music on fetalbehavior and various possible biological, psychological, and social mechanisms are discussed. The potential preventive effect of music-based interventions, approaches, and practices and their possible use across different cultures are also considered. Overall, we highlight how music, employed in a variety of ways, may support perinatal mental health with the aim of stimulating more interest and research in this area.

Magdalena et al. (2021) conducted a study on ‘Improvement of the breastfeeding self-efficacy and postpartum mental health after lactation consultations –An Observational study’. The results suggest that women willing to benefit from the lactation consultations may exhibit symptoms of mental health difficulties that may be associated with difficulties in breastfeeding. Strengthening breastfeeding confidence during lactation consultations may reflect in the improvement of the woman's mental health.

Recommendations

Education and Awareness

Informing expectant mothers about the typical emotional changes that occur during pregnancy and the support options that are available can lessen worry.

Counseling and Therapy

Offering counseling or therapy to people who are depressed or anxious before having a baby might give them coping mechanisms and emotional support.

Social Support

Social support can help pregnant women feel less alone and stressed by encouraging them to interact with friends, family, or support groups.

Mindfulness and Relaxation Techniques

Teaching mindfulness and relaxation techniques, such as yoga or meditation, can aid in managing stress and anxiety.

Community Programmes

Putting in place community-based initiatives that give expecting moms access to resources and assistance for their mental health.

Recommendations on Postnatal Period and Women's Mental Health

Mother-Baby Bonding Support

Offering advice on attachment and bonding methods to strengthen the emotional relationship between mother and child.

Partner Involvement

The stress on the new mother might be lessened by encouraging partners to take an active role in providing care and emotional support.

Postnatal Support Groups

Establishing secure settings where new mothers can talk about their experiences and get emotional support from other women.

Peer Mentoring Programmes

Programmes that pair new mothers with seasoned mothers who have faced comparable obstacles can offer invaluable advice and compassion.

These suggestions are meant to develop an extensive system of support that attends to the needs of new and expecting mothers holistically, from early identification and intervention to community and workplace support. By putting these strategies into practice, we may make great progress toward enhancing mothers' mental health and, in turn, the well-being and growth of their offspring and families.

Future Implications:

Mental health during pregnancy and after delivery is a serious issue, and its importance will only increase. It is crucial to acknowledge and address the mental health of expecting and new moms since it affects not only the people who are directly impacted but also families, society, and the environment.

The effects of maternal mental health on the family are extensive. The mother's emotional state during pregnancy can affect fetal development and the future mental health of the unborn child. Postpartum maternal mental discomfort can affect the child's emotional and cognitive development by impairing the mother's ability to bond with the infant and providing insufficient care. Due to their role as carers and potential stress, partners and other family members are also impacted. The mental health of moms is crucial for healthy family dynamics and child development.

The implications are significant from a societal perspective. Untreated maternal mental health problems have a cost in higher medical costs, decreased productivity, and higher rates of child abuse. The workforce is impacted because mothers may find it difficult to find

employment, which limits their financial potential. This, therefore, has an effect on the whole economy. In addition, untreated maternal mental health problems might cause mental disease relapses that will damage future generations.

In terms of the environment, the ripple effects are subtle but considerable. A family dealing with maternal mental health concerns might spend more money on support and care. Additionally, untreated mental health issues might encourage excessive consumption and wasteful behavior because people may resort to negative coping techniques.

Determining how to address antenatal and postnatal mental health is crucial. Destigmatization, readily available mental health care, and early intervention are essential. In addition to creating healthier families, promoting maternal mental health also benefits a more prosperous society and a more sustainable environment. It's a comprehensive strategy that acknowledges the interdependence of societal, familial, human, and environmental well-being, elevating it to a top priority for the future.

Conclusion

Essentially, it is impossible to stress the importance of giving antenatal and postnatal mental health a top priority. Pregnancy and childbirth are transforming stage of life that includes both joys and difficulties. It is clear that maintaining mothers' mental health throughout these times is crucial for the overall well-being of mothers, children, and families.

The emotional complexity that expecting and new moms go through highlights the need for specialized support. Neglecting mental health throughout pregnancy and after delivery can have negative effects on both the mother's and the baby's well-being. According to research, there is a clear connection between a mother's mental health and her child's development, and this connection has a long-lasting effect on a child's emotional and cognitive development.

An all-encompassing strategy is needed to address the mental health of expectant and new mothers. In order to ensure that women receive timely help, healthcare organizations should incorporate mental health exams into normal care. Peer support groups and community organizations working together to provide safe venues for women to discuss their stories helps lessen feelings of loneliness.

In addition, how maternal mental health is treated is greatly influenced by political regulations. It is crucial to allocate funds for antenatal and postnatal women's mental health treatments. A comprehensive strategy that recognizes the special problems encountered by new mother’s benefits from policies that provide flexible maternity leave and encouraging work settings.

In conclusion, it is critical to acknowledge and treat the importance of antenatal and postnatal mental health. Mothers' mental health should be prioritized since it leads to healthier families, more robust communities, and a more adaptable society. For the benefit of both the present and future generations, now is the moment to take action.

References

Argent, V., Sherratt, M., & Nelson, M. (2022). A Perinatal Mental Health Education Programme for Maternity, Neonatal and Other Allied Multidisciplinary Teams in NHS Lithian: Why, How, and What's Next. BJPsych Open, 8(S1), S14-S15.

Barkin, J. L., Philipsborn, R. P., Curry, C. L., Upadhyay, S., Geller, P. A., Pardon, M., ... &Buoli, M. (2022). Climate change is an emerging threat to perinatal mental health. Journal of the American Psychiatric Nurses Association, 10783903221139831.

Chrzan-Dętkoś, M., Walczak-Kozłowska, T., Pietkiewicz, A., &Żołnowska, J. (2021). Improvement of the breastfeeding self-efficacy and postpartum mental health after lactation consultations–Observational study. Midwifery, 94, 102905.

Clayborne, Z. M., Colman, I., Kingsbury, M., Torvik, F. A., Gustavson, K., & Nilsen, W.(2022). Prenatal work stress is associated with prenatal and postnatal depression and anxiety: Findings from the Norwegian Mother, Father and Child Cohort Study (MoBa). Journal of affective disorders, 298, 548-554.

Endres, K., Haigler, K., Sbrilli, M., Jasani, S., & Laurent, H. (2023). Social determinants of perinatal mental health during the COVID-19 pandemic. General Hospital Psychiatry, 84, 3943.

Heuckendorff, S., Johansen, M. N., Johnsen, S. P., Overgaard, C., & Fonager, K. (2021). Parental mental health conditions and use of healthcare services in children the first year of life–a register-based, nationwide study. BMC Public Health, 21(1), 1-12.

Key nejad, R. C., Bitew, T., Mulushoa, A., Howard, L. M., & Hanlon, C. (2023). Pregnant women and health workers' perspectives on perinatal mental health and intimate partner violence in rural Ethiopia: a qualitative interview study. BMC pregnancy and childbirth, 23(1), 78.

Kokkinaki, T., &Hatzidaki, E. (2022). COVID-19 pandemic-related restrictions: Factors that may affect perinatal maternal mental health and implications for infant development. Frontiers in Pediatrics, 10, 846627.

Lever Taylor, B., Kandiah, A., Johnson, S., Howard, L. M., & Morant, N. (2021). A qualitative investigation of models of community mental health care for women with perinatal mental health problems. Journal of Mental Health, 30(5), 594-600.

Mokhtari Zanjani, P., Tehranian, N., Changizi, N., Mohammadi, E., Farzinrad, B.,&Kazemnejad, A. (2023). Maternal mental health concerns during the COVID-19 pandemic in Iran: a content analysis study. Reproductive Health, 20(1), 102.

Ryan, A., & Barber, C. (2022). Postnatal depression and anxiety during the COVID-19 pandemic: The needs and experiences of New Zealand mothers and health care providers. Midwifery, 115, 103491.

Sanfilippo, K. R. M., Stewart, L., & Glover, V. (2021). How music may support perinatal mental health: an overview. Archives of women's mental health, 24(5), 831-839.

Schaming, C., & Wendland, J. (2022). Postnatal mental health during the COVID-19 pandemic: Impact on mothers' postnatal sense of security and on mother-to-infant bonding. Midwifery, 117, 103557.

Varma, D. S., Mualem, M., Goodin, A., Gurka, K. K., Wen, T. S. T., Gurka, M. J., & Roussos-Ross, K. (2023). Acceptability of an mHealth App for Monitoring Perinatal and Postpartum Mental Health: Qualitative Study With Women and Providers. JMIR Formative Research, 7, e44500.

Wilkinson, K. (2023). Maternal (perinatal) mental health and employment: An agenda for research and practice. Human Resource Management Journal, 33(2), 346-361.